

# Reservoir Sediment Management & Analysis for Engineers


## Downstream Channel Impacts of Reservoir Sediment Management

**John Shelley, Ph.D., P.E.**  
Kansas City District

University of Kansas  
LEEP2 Building – Room G415  
June 11-15, 2018



**ERDC**  
Engineer Research and  
Development Center



# Outline


- Natural rivers
- Effects of drawdown flushing
- Effects of sediment bypass



# Impacts from Lack of Turbidity: Colorado River

David Ward and Rylan Morton  
Starnes, USGS, Grand Canyon  
Monitoring and Research Station

- Humpback Chub numbers have decreased substantially and they are now federally protected
- One primary reason is that the Colorado River used to be usually over 1000 FNU, but after construction of Glen Canyon Dam now is usually below 50 FNU. The small chub become easy prey for trout species in clear water.




**Brown trout**  
**mean TL = 261 mm**



**Humpback chub**  
**mean TL = 56 mm**

# Unregulated Rivers Experience High Sediment Loads During Floods





Missouri River at Sioux City, IA. Data is approximate for demonstration purposes only.



# Downstream Impacts Depend On Two Major Factors

1-How closely does the management option match the natural timing, concentration, and gradation of incoming sediment load?



2-What the downstream channel (and users) are “used to”

- ▶ Historically-turbid Midwest stream vs. Historically-clear mountain stream
- ▶ When was the downstream channel infrastructure built (and for what conditions?)
- ▶ Sport fish vs. native fish?

# Outline

- ✓ ■ Natural rivers
  - Effects of drawdown flushing
  - Effects of sediment bypass



# Effects of Drawdown Flushing

## US Examples

- Willwood
- Spencer Dam
- Fall Creek

Minimizing Downstream Channel Impacts





The Willwood Diversion Dam was built in 1924 and still contains some original equipment that could only be replaced by drawing the water behind the dam down.

[http://billingsgazette.com/lifestyles/recreation/muddy-water-suffocates-fish-in-prime-stretch-of-wyoming-s/article\\_94ad366d-a43b-542b-9703-b832b3900606.html](http://billingsgazette.com/lifestyles/recreation/muddy-water-suffocates-fish-in-prime-stretch-of-wyoming-s/article_94ad366d-a43b-542b-9703-b832b3900606.html)





+10



[billingsgazette.com](http://billingsgazette.com)

Piles of sediment have built up behind the Willwood Diversion Dam, which was built in 1924.



+10

[billingsgazette.com](http://billingsgazette.com)

This view looks downstream from the dam.







[billingsgazette.com](http://billingsgazette.com)





# Spencer Dam Flushing





# Spencer Dam Flush- The Next Morning



# Spencer Dam Fish Kills- 1977-1979

Hess and Newcomb (1982)  
document fish kills

DO dropped to 3.1 mg/L

Species	Number
Channel catfish ( <i>Ictalurus punctatus</i> )	954
Walleye ( <i>Stizostedion vitreum</i> )	9
Sauger ( <i>Stizostedion canadense</i> )	94
Green sunfish ( <i>Lepomis cyanellus</i> )	31
Bullhead ( <i>Ictalurus</i> spp.)	10
Madtom ( <i>Noturus gyrinus</i> )	4
Bluegill ( <i>Lepomis macrochirus</i> )	29
Gizzard shad ( <i>Dorosoma cepedianum</i> )	2
Largemouth bass ( <i>Micropterus salmoides</i> )	78
Drum ( <i>Aplodinotus grunniens</i> )	158
Crappie ( <i>Pomoxis</i> spp.)	41
Common carp ( <i>Cyprinus carpio</i> )	560
Shorthead redhorse ( <i>Moxostoma macrolepidotum</i> )	763
River carpsucker ( <i>Carpionodes carpio</i> )	819
Stonecat ( <i>Noturus flavus</i> )	58
Goldeye ( <i>Hiodon alosoides</i> )	2
Northern pike ( <i>Esox lucius</i> )	5
Brown trout ( <i>Salmo trutta</i> )	4
Smallmouth bass ( <i>Micropterus dolomieu</i> )	2
White sucker ( <i>Catostomus commersoni</i> )	2
Minnow (Cyprinidae)	18,846
Total	22,471



# Spencer Dam: Operational Changes since 1989

The pool is drained slowly before the flushing begins.

“The operational modifications of raising the gates slowly and dropping the hydro pond at a reduced rate has been successful in avoiding fish kills since 1989.”

--Gutzmer, King, and Overhue 1996

“It appears that with environmentally friendly ways to pass sediment, fish below Spencer Dam survive and express resilience to conditions created by sluicing.”

--Gutzmer, King, Overhue, and Chrisp 2002



Why would lowering the pool  
slowly before opening the  
flushing gates reduce fish kills?





# Fall Creek Lake Drawdown Flush







Photo Source: Greg Taylor



# Suspended Sediment Loads


## Fall Creek Outflow – 2012/13



- Pre-drawdown: 4,300 tons (34 days)
- Drawdown: 51,600 tons (6 days)
- Post-drawdown: 4,030 tons (53 days)







Schenk and Bragg (2014)


**Figure 9.** Dissolved oxygen, turbidity, and streamflow during the drawdown at *Fall Creek Outflow*, Middle Fork Willamette River Basin, Oregon, December 2012.


“During the drawdown, DO data at Fall Creek Outflow decreased rapidly at the onset of the large sediment release from approximately 13 mg/L to a minimum value of 6.50 mg/L”



# Dissolved Oxygen and Turbidity, November 2015

Fall Creek below Winberry Creek, near Fall Creek, OR (14151000)





“DO at Jasper decreased slightly during the drawdown to a minimum value of 11.63 mg/L, suggesting that although a small effect is possible, the sediment release from Fall Creek Lake did not cause a rapid decrease in DO approximately 10 river miles downstream of the dam.” --Schenk and Bragg (2014)



# Fall Creek Sediment Flushing



- Ten-fold increase in the adult salmon that later return to Fall Creek
- No observed fish kills
- In-reservoir: Removal of invasive species, significant increase in natural populations

# Effects of Drawdown Flushing

## US Examples

- Willwood
- Spencer Dam
- Fall Creek

Minimizing Downstream Channel Impacts





# Minimizing Downstream Impacts

- Mimic natural conditions
  - ▶ Max SSC = flood SSC
  - ▶ Time of year = natural flooding time of year
- Minimize fish impacts
  - ▶ Assess with Severity Index
  - ▶ Dilute sediment discharges
  - ▶ Alternate clear water and sediment laden discharges



# Newcomb and Jenson (1996) Meta Analysis

- 80 studies
- Six simple, empirical equations relating severity of ill effects on fish to
  - ▶ SSC in mg/l
  - ▶ Duration in hours



SEV	Description of effect
<b>Nil effect</b>	
0	No behavioral effects
<b>Behavioral effects</b>	
1	Alarm reaction
2	Abandonment of cover
3	Avoidance response
<b>Sublethal effects</b>	
4	Short-term reduction in feeding rates; short-term reduction in feeding success
5	Minor physiological stress; increase in rate of coughing; increased respiration rate
6	Moderate physiological stress
7	Moderate habitat degradation; impaired homing
8	Indications of major physiological stress; long-term reduction in feeding rate; long-term reduction in feeding success; poor condition
<b>Lethal and para-lethal effects</b>	
9	Reduced growth rate; delayed hatching; reduced fish density
10	0–20% mortality; increased predation; moderate to severe habitat degradation
11	>20–40% mortality
12	>40–60% mortality
13	>60–80% mortality
14	>80–100% mortality

# Juvenile and Adult Salmonids

Duration of exposure to SS ( $\log_e$  hours)

0	1	2	3	4	5	6	7	8	9	10
---	---	---	---	---	---	---	---	---	---	----

(A) Average severity-of-ill-effect scores (empirical)

162755	14	-	-	14	-	-	-	-	-	-	-	12
59874	-	-	12	10	14	12	-	-	-	-	-	11
22026	-	-	-	-	-	11	-	-	-	-	-	10
8103	3	-	-	10	12	10	-	11	-	14	-	9
2981	6	-	-	9	11	8	9	13	-	-	-	8
1097	4	-	-	-	10	9	11	8	-	14	8	7
403	4	5	8	-	-	8	10	9	10	-	-	6
148	4	-	6	9	-	8	9	9	-	-	-	5
55	2	4	5	-	-	7	10	9	10	-	-	4
20	3	4	-	4	-	-	-	10	-	9	-	3
7	-	-	-	-	-	3	-	9	-	9	-	2
3	-	-	-	-	-	5	-	-	-	-	-	1
1	3	-	-	-	-	-	-	-	-	-	-	0
	1	3	7	1	2	6	2	7	4	11	30	
	Hours			Days			Weeks		Months			



# Adult Freshwater Nonsalmonids

Duration of exposure to SS ( $\log_{10}$  hours)

0	1	2	3	4	5	6	7	8	9	10
---	---	---	---	---	---	---	---	---	---	----

(A) Average severity-of-ill-effect scores (empirical)

Concentration (mg SS/L)

162755	10	-	-	-	-	-	10	-	-	-	-	12
59874	-	-	-	-	-	-	-	-	-	-	-	11
22026	-	-	-	-	-	-	10	-	-	-	-	10
8103	5	-	-	-	-	-	-	-	-	-	-	9
2981	-	-	-	-	-	-	-	-	-	13	-	8
1097	-	-	-	-	-	-	-	12	-	-	-	7
403	4	-	-	-	10	-	-	-	-	-	-	6
148	-	-	-	-	-	-	10	11	-	-	-	5
55	-	-	-	-	-	-	-	9	-	-	-	4
20	4	-	-	-	-	-	-	-	-	12	-	3
7	-	-	-	-	-	-	-	-	-	-	-	2
3	-	-	-	-	-	-	-	-	-	-	-	1
1	-	-	-	-	-	-	-	-	-	-	-	0
	1	3	7	1	2	6	2	7	4	11	30	
	Hours			Days			Weeks		Months			

( $\log_{10}$  mg SS/L)

# Adult Freshwater Nonsalmonids

Duration of exposure to SS ( $\log_e$  hours)

0	1	2	3	4	5	6	7	8	9	10
---	---	---	---	---	---	---	---	---	---	----

(B) Average severity-of-ill-effect scores (calculated)

Concentration (mg SS/L)	7	8	9	10	10	11	12	12	13	14	-	12
	7	8	9	9	10	11	11	12	13	14	14	11
	7	8	8	9	10	10	11	12	13	13	14	10
	7	7	8	9	9	10	11	12	12	13	14	9
	6	7	8	8	9	10	11	11	12	13	13	8
	6	7	7	8	9	10	10	11	12	12	13	7
	6	6	7	8	9	9	10	11	11	12	13	6
	5	6	7	8	8	9	10	10	11	12	13	5
	5	6	7	7	8	9	9	10	11	12	12	4
	5	6	6	7	8	8	9	10	11	11	12	3
	5	5	6	7	7	8	9	10	10	11	12	2
	4	5	6	7	7	8	9	9	10	11	11	1
	4	5	6	6	7	8	8	9	10	10	11	0
1			3	7	1	2	6	2	7	4	11	30
Hours			Days			Weeks			Months			

( $\log_e$  mg SS/L)

# For more information:

Newcombe, C. and Jensen, J. 1996.


“Channel Suspended Sediment and Fisheries: A Synthesis for Quantitative Assessment of Risk and Impact”

North American Journal of Fisheries Management, Volume 16, November 1996, Number 4.





# Lake Como (Italy)



# Lake Como (Italy)

- Non-consecutive days
  - ▶ 2009: 16 days from 23 May to 10 July
  - ▶ 2010: 6 days from 8 July to 20 July
  - ▶ 2011: No flush
- SSC Thresholds:
  - ▶ 1,500 mg/L daily average
  - ▶ 3,000 mg/L alert value to adjust ongoing activity



# Lake Como (Italy)



- 44% fines
- 54% sand



# Results

- Macroinvertebrates
  - ▶ Seasonal changes, no change during flush years vs. non flush year
- Bullhead (EU protected)
  - ▶ Increase in density
- Brown trout
  - ▶ No impact





# Outline

- ✓ ■ Natural rivers
- ✓ ■ Effects of drawdown flushing
  - Effects of sediment bypass



# Two types of bypass (as far as the downstream channel is concerned)



1-Options that raise the base-level SSC

2-Options that raise the flood-related SSC

If “raise” = “restore” the effect is generally positive from an ecological perspective, though there could still be infrastructure concerns.







# Millsite Lake



# Matching Incoming Sediment




# Solis Reservoir, Switzerland






# Solis Reservoir, Switzerland







Oertli and Auel, 2015





# Downstream Discharge




Oertli and Auel, 2015



# Solis Reservoir, Switzerland

## Measured

- Chemical properties
- Sediment respiration
- Benthic organic matter
- Sediment size distribution
- Periphyton and macroinvertebrates



5x/year for two years



# Solis Reservoir, Switzerland:

## Conclusions

- Sediment Bypass Tunnel (SBT) discharges induced effects in the downstream channel typical of natural flooding.
- “In the short term, SBT operations can increase the flow/sediment variability that is often lost in flow-regulated rivers.”
- “A permanent positive change in the system would take several years of adaptive management operations, similar to experimental floods.”
- “In conclusion, we found that SBTs, if ecologically implemented (i.e. having the operational characteristics similar to the natural flood features of a system), can improve the longitudinal connectivity of sediments of rivers impounded by dams. Indeed, SBT events can be used as environmental flows to simulate more natural flow/sediment regimes of receiving waters.”



Martin, Doering, and Robinson, 2017



# Conclusion

- ✓ ■ Natural rivers
  - ▶ How closely does the sediment discharge match the natural, no-dam conditions?
- ✓ ■ Effects of drawdown flushing
  - ▶ How can the sediment concentrations be limited to acceptable severity levels?
- ✓ ■ Effects of sediment bypass





# However...

- You will have to work closely with your state environmental agencies.
- They may need to become accustomed to the idea that sediment is a resource not a pollutant.



- We put on a workshop specifically for regulators last year.
  - ▶ Highly successful at changing paradigms.
- We would be happy to tailor a workshop to your state agencies and hold it in your area.

